

SENSORY REWIRING IN AN ECHOLOCATOR: GENOME-WIDE MODIFICATION OF
RETINOGENIC AND AUDITORY GENES IN THE BAT *MYOTIS DAVIDII*

Nicholas J. Hudson^{1*}, Baker ML², Hart NS³, Wynne JW², Gu Q^{4,5}, Huang Z⁶, Zhang G⁶, Ingham AB¹,
Wang L-F^{2,7} and Reverter A¹

¹ Computational and Systems Biology, CSIRO Agriculture Flagship, Queensland Bioscience Precinct,
306 Carmody Road, Brisbane, Queensland, Australia

² Australian Animal Health Laboratory, CSIRO Biosecurity Flagship, Geelong, Victoria, Australia

³ School of Animal Biology and the Oceans Institute, University of Western Australia, Crawley, WA
6009, Western Australia, Australia

⁴ College of Information Sciences and Technology, Donghua University, Shanghai 201620, China

⁵ Faculty of Medicine, Imperial College London, South Kensington, London SW7 2AZ, England

⁶ BGI-Shenzhen, Shenzhen, 518083, China

⁷ Program in Emerging Infectious Diseases, Duke-NUS Graduate Medical School, Singapore

*Author for correspondence

Dr. Nicholas James Hudson

nick.hudson@csiro.au

DOI: 10.1534/g3.114.011262

Files S1-S2

Available for download as Excel files at <http://www.g3journal.org/lookup/suppl/doi:10.1534/g3.114.011262/-/DC1>

File S1 CUB analysis for *M. davidii* and *P. alecto*

File S2 CUB analysis for *M. lucifugus* and *P. vampyrus*

Figure displaying BioGPS normalized tissue expression for the gene TSPAN10, highly biased in *M. lucifugus*.

