

Table S1 Oligonucleotide primers used for PCR.

Translocation	Breakpoint		
	A	B	C
EB4	GCCGGTTTTGGAGCA	GCGGGCGGCAAA	TCCACACCAGAG
	TCCATACACAGGG	GGCTGTT	GTCGTAG
	AAGTTGTAGCTACGC	GGGCGGCTTCGGC	ATCAATTCGCGA
	TGAAACACCAGATG ACC	AGTAAA	TGGAACGG
IBj5	CTCTCGCCCGACTAG	GCTGTGACTCATA	
	GACTTC	CTTCCCC	-
	GTTGCCCTGCTTTCC	GTTCGCTAGTGAG	
	GTGCG	TGCGTTCC	
UK14-1	GGTAGGTAAGGAAG		
	GTGCAATCG	-	NA
	CGATGAAGAGAGGC		
	CCAGTGAAGAC		
B362i	ATAGTGGGAGCTGTC	AGCTCGAATCGCG	TTCATCGAGACC
	ACAGGTTCCCTTG	AGGAGAG	GGCTGGAAG
	AAGTTGTAGCTACGC	GTCTTCGGGCTTC	CGCGATGTCACC
	TGAAACACCAGATG ACC	AACCGAG	GACGAAAG

NA – not applicable.

Oligonucleotide primers used to PCR amplify *ad-7* sequences of *RIP3C* and *RIP3T*

RIP3C CTCCAACCTAGCACT
 TGTTGATC

 GGTTGAGGCTGATAT
 CCATTCTCC

RIP3T GTATGTATCCGTCCC
 CTCTCATCTC

 GAAGCCAAAGCAAT
 CGGTCG
